

Nutrition 101

Jason Banta, Ph.D., PAS
 Asst. Professor and Extension Beef Cattle Specialist
 Texas A&M AgriLife Extension Service
 Texas A&M University
 Overton, TX

1

6 major nutrients groups

- proteins
- carbohydrates
- lipids
- vitamins
- minerals
- water



2

Nutrients Cattle Require

- | | |
|---|--|
| <ul style="list-style-type: none"> • protein • energy <ul style="list-style-type: none"> - carbohydrates - lipids <ul style="list-style-type: none"> · short or long chain - excess protein • vitamins <ul style="list-style-type: none"> - A, D, E, K - B-vitamins | <ul style="list-style-type: none"> • minerals <ul style="list-style-type: none"> - macro <ul style="list-style-type: none"> · Ca, P, K, Mg, Na, S - micro <ul style="list-style-type: none"> · Cu, Zn, Mn, Se, I, Co, Fe, • water • other nutrients <ul style="list-style-type: none"> - linoleic acid - linolenic acid - etc. |
|---|--|

*nutrients in red are not essential in the diet
 (i.e. they can be supplied by rumen microbes)

3

Nutrient Usage

- | | |
|---|---|
| <ul style="list-style-type: none"> • maintenance • pregnancy • lactation • gain | <ul style="list-style-type: none"> energy protein minerals vitamins water all other nutrients |
|---|---|

maintenance energy – the amount of energy it takes to maintain an animal (i.e. the animal is not gaining or losing weight or condition)

4

total digestible nutrients (TDN)

> TDN = energy

crude protein (CP)

5

highest nutrient needs as % of diet DM

calves

stocker cattle

lactating cows

replacement heifers

dry cows

lowest nutrient needs



6

Description	% CP	% TDN	% Ca	% P
2-yr-old lactating cow, peak lactation	11.5	60	0.28	0.18
3-yr-old lactating cow, peak lactation	12.5	61	0.30	0.19
mature lactating cow, peak lactation	12.5	61	0.30	0.19
coming 3-yr-old dry cow, 270 d pregnant	9.0	58	0.26	0.17
mature dry cow, 270 d pregnant	8.5	55	0.26	0.17

*Estimated dietary requirements for high marbling cows with no weather stress. Assumes 1,300 lb mature weight and 25 lb milk potential at maturity (NRC, 2016)

7

Nutrient Requirements

700 lb yearling steer

ADG	% TDN	% CP	% Ca	Ca, gm	DMI, lb
1.0	53	8	0.26	21.3	18.1
1.5	57	9	0.32	26.5	18.5
2.0	61	10	0.38	31.6	18.7
2.5	65	11	0.43	36.1	18.7
3.0	70	12	0.50	41.7	18.7

*Estimated dietary requirements for Brangus type steer under typical production conditions (Beef Cattle NRC, 1996). These requirements will vary depending on numerous factors including body condition, health, breed, environmental factors, use of growth promotants, and others.

8

Nutrient Requirements

comparison at 2.0 lb/d


weight	% TDN	% CP	% Ca	Ca, gm	DMI, lb
500	65	12.7	0.55	32.2	13.0
600	63	11	0.45	32.3	15.9
700	61	10	0.38	31.6	18.7
800	61	9.5	0.34	31.4	20.6

*Estimated dietary requirements for Brangus type steer under typical production conditions (Beef Cattle NRC, 1996). These requirements will vary depending on numerous factors including body condition, health, breed, environmental factors, use of growth promotants, and others.


9



10



11



12

Acidosis

13



pH 0	Battery Acid
pH 1	Stomach Acid
pH 2	Lemon Juice, Vinegar
pH 3	Orange Juice, Soda, Some Dental Rinses
pH 4	Tomato Juice, Beer
pH 5	Black Coffee
pH 6	Saliva, Cow's Milk
pH 7	Pure Water
pH 8	Sea Water, pH-Neutralizing Dental Rinses
pH 9	Baking Soda
pH 10	Antacids
pH 11	Antacids, Dental Treatment Rinses
pH 12	Soapy Water

14



(photo courtesy of vetnext.com)

15



(photo courtesy of FiveF)

16



17



18

TMR VS. Supplementation

19

When do we have confined feeding?

- feedlot
- dairy
- backgrounding yard
- drought
- winter in Northern states

20

TMR

3 components

- roughage
- energy source
- protein, mineral, vitamin premix

21

When do we supplement?

- forage doesn't meet nutrient requirements
- BCS is too low
- improve forage utilization
- improve ADG of growing cattle

22

Energy and/or Protein Supplements

components

- not worried about roughage
- energy
- protein

separate mineral & vitamin supplement
generally provided

23


Performance Terminology

Growing Cattle: ADG


Cows: BCS




24




25




26



27




28




29



30




31



32



33




34




35



36



37




38

When should you take BCS?

- at weaning
 - most important time
- every few months
- score the same cows in the herd
- a digital camera can be a good tool


39




40

BCS and Pregnancy Rates

41



42




43


<http://beef.tamu.edu>

jpbanta@ag.tamu.edu


44




7



8



9



10

Cow-Calf Systems are based on Perennial Forages

11

Native Warm-Season Perennial Forages

12



13

(photo courtesy of Steve Hammack)



14

(photo courtesy of Bruce Carpenter)

Protein is often
first limiting with
Native Warm-Season
Perennial Forages

15

Native Range Systems

16



17

Introduced
Warm-Season
Perennial Forages

18



19



20



(photo courtesy of Joe Paschal)

21



(photo courtesy of Joe Paschal)

22



(photo courtesy of Joe Paschal)

23

Energy is often first limiting with Introduced Warm-Season Perennial Forages

24


Bahia and Bermudagrass Systems

25

How long does bahia and bermudagrass grow?

26

Utilizing Warm-Season Perennial Grasses (bermudagrass, bahiagrass, etc.)



27

How do we fill in the gaps?

28



29



30



31

What about winter annuals?

32




33



34



35



36



37



38

What about
summer annuals?

39



40



41

Management Factors
Affecting
Forage Production

42



43

Grazing Pressure and Stocking Rate

44



45



46



47



48



49




50




51



52



53



54

Stocking Strategies and Factors


55

Stocking Rate is a Moving Target

56

How many acres do you need per cow?

57



58

general intake guidelines

- dry gestating cow: 1.8 to 2.0% of BW
- lactating cow: 2.3 to 2.5% of BW



(intake guidelines from Lardy et al., 2004)

59

Estimated Annual Intake

assume cow consumes 2.25% of BW on DM basis

$$1,000 \text{ lb} \times 2.25\% = 22.5 \text{ lbs/d}$$

$$22.5 \times 365 = \underline{8,213 \text{ lbs/year}}$$

60

Estimated Forage Allowance

Should you let the cows consume everything that is produced?

let cow consume 70%

$$8,213 \div 0.70 = \underline{11,733 \text{ lbs}}$$

61



62

70% consumption

Cow Weight	1,000	1,200	1,400
Cow intake, (2.25% of BW)	8,213	9,855	11,498
Calf Intake	1,508	1,810	2,111
Pair Intake	9,721	11,665	13,609
Forage Allowance, let cow consume (70%)	13,887	16,664	19,441
Forage Production, lbs/ac	4,500	4,500	4,500
Stocking Rate, ac/pair	3.09	3.70	4.32

63




(photo courtesy of Bruce Carpenter)

64

25% consumption

Cow Weight	1,000	1,200	1,400
Cow intake, (2.25% of BW)	8,213	9,855	11,498
Calf Intake, lbs/year	1,508	1,810	2,111
Pair Intake, lbs/year	9,721	11,665	13,609
Forage Allowance, let cow consume (25%)	38,884	46,660	54,436
Forage Production, lbs/ac/yr	3,000	3,000	3,000
Stocking Rate, ac/pair	12.96	15.55	18.15

65



66

Using Excess Forage

67



68



skip an application of nitrogen

69




70



71



72




be mindful of biosecurity

73



be mindful of biosecurity

74



be mindful of biosecurity

75



beef.tamu.edu

jpbanta@ag.tamu.edu

76